



Fracture Liaison Service Database

Annual report 13 January 2022

References and bibliography

Black DM, Arden NK, Palermo L, Pearson J, Cummings SR. Prevalent vertebral deformities predict hip fractures and new vertebral deformities but not wrist fractures. Study of Osteoporotic Fractures Research Group. *J Bone Miner Res.* 1999;14:821-28.

Howlett, D, Drinkwater, K, Mahmood, N, Illes, J, Griffin, J and Javaid, K. Radiology reporting of osteoporotic vertebral fragility fractures on computed tomography studies: results of a UK national audit. *European Radiology.* 2020;30:4713-23.

National Institute for Health and Care Excellence. Bisphosphonates for treating osteoporosis. NICE technology appraisal guidance (TA464). Manchester: NICE, 2017. www.nice.org.uk/guidance/ta464

National Institute for Health and Care Excellence. Denosumab for the prevention of osteoporotic fractures in postmenopausal women. NICE technology appraisal guidance (TA204). Manchester: NICE, 2010. www.nice.org.uk/guidance/ta204

National Institute for Health and Care Excellence. Falls in older people: assessing risk and prevention. NICE clinical guideline (CG161). Manchester: NICE, 2013. www.nice.org.uk/guidance/cg161

National Institute for Health and Care Excellence. Falls in older people. NICE clinical guideline (QS86). Manchester: NICE, 2015. <https://www.nice.org.uk/guidance/qs86>

National Institute for Health and Care Excellence. Osteoporosis: assessing the risk of fragility fracture. NICE clinical guideline (CG146). Manchester: NICE, 2012. <https://www.nice.org.uk/guidance/cg146>

National Institute for Health and Care Excellence. Osteoporosis. NICE quality standard (QS149). Manchester: NICE, 2017. www.nice.org.uk/guidance/qs149

National Institute for Health and Care Excellence. Raloxifene and teriparatide for the secondary prevention of osteoporotic fragility fractures in postmenopausal women. NICE clinical guideline (TA161). Manchester: NICE, 2008. <https://www.nice.org.uk/guidance/ta161>

National Osteoporosis Guideline Group. NOGG 2017: Clinical guideline for the prevention and treatment of osteoporosis. Sheffield: NOGG, 2017. www.sheffield.ac.uk/NOGG/

O'Neill TW, Felsenberg D, Varlow J, Cooper C, Kanis JA, Silman J. The prevalence of vertebral deformity in European men and women: the European Vertebral Osteoporosis Study. *J Bone Miner Res.* 1996;11:1010-18.



Royal College
of Physicians

FLS-DB

Royal College of Physicians
11 St Andrews Place, Regent's Park
London NW1 4LE

+44 (0)20 3075 1511
flsdb@rcp.ac.uk
rcp.ac.uk

Royal College of Physicians. Fracture Liaison Service Database annual report. Benchmarking FLS improvement and performance in 2019: Pre-COVID. London: RCP, 2021.

Royal Osteoporosis Society. Clinical guidance for the effective identification of vertebral fractures. Bath: ROS, 2017. <https://theros.org.uk/media/3daohfrq/ros-vertebral-fracture-guidelines-november-2017.pdf>

Royal Osteoporosis Society. Effective secondary prevention of fragility fractures: clinical standards for fracture liaison services. Bath: ROS, 2015. <https://theros.org.uk/media/1776/clinical-standards-report.pdf>

Royal Osteoporosis Society. Quality Standards for Osteoporosis and Prevention of Fragility Fractures. Bath: ROS, 2017. <https://theros.org.uk/media/99099/op-standards.pdf>